

EFFECT OF KILLING AND TECHNOLOGICAL PROCESSING ON MICROBIOLOGICAL QUALITY OF EDIBLE INSECTS

Petra Škvorová*¹, Martin Kulma², Lenka Kouřimská¹, Roman Švejstl¹

Address(es): Ing. Petra Škvorová,

¹ Department of Microbiology, Nutrition and Dietetics, Czech University of Life Sciences, Prague, Kamýcká 129, 165 00, Praha-Suchdol, Czech Republic.

² Department of Zoology and Fisheries, Czech University of Life Sciences, Prague, Kamýcká 129, 165 00, Praha-Suchdol, Czech Republic.

*Corresponding author: skvorova@af.czu.cz

<https://doi.org/10.55251/jmbfs.11018>

ARTICLE INFO

Received 9. 2. 2024
Revised 30. 7. 2024
Accepted 1. 7. 2024
Published 1. 8. 2024

Regular article



ABSTRACT

This study aimed to assess the effect of killing (blanching and freezing) followed by culinary processing on the microbiological quality of *Tenebrio molitor* and *Gryllus assimilis*. These insects were subjected to starvation before killing, which does not ensure sufficient microbiological safety. Therefore, technological or culinary processing of insects is necessary before consumption. For the subsequent processing of insects, boiling, drying, roasting, and microwave treatment were used. Nine categories of microbiological parameters were selected as indicators of hygienic quality. All identification and enumeration were performed using ISO-compliant methods. The results showed that killing insects by blanching or freezing alone does not ensure sufficient microbiological quality, and therefore, further culinary processing is necessary. However, all subsequent culinary treatments applied in this study were able to reduce the microbial load significantly.

Keywords: *Tenebrio molitor*, *Gryllus assimilis*, roasting, boiling, microwave treatment, drying

INTRODUCTION

Due to the increasing population, meat production and consumption are expected to reach unsustainable levels in the coming decades. Therefore, new and sustainable sources of nutrients, especially protein, must be sought (van Huis and Ooninx, 2017). The urgency in finding alternative and sustainable protein sources has triggered an exponential increase in interest in insects as a source of human food or livestock feed over the past decade (van Huis et al., 2021; Shah et al., 2022).

Compared to conventional livestock, insects have similar nutritional value and a lower environmental impact. However, unlike conventional livestock, their chemical composition is influenced by various factors, including species, developmental stage, sex, diet, rearing and processing methods (Ooninx and Finke, 2021). Generally, proteins and lipids, the major nutrients in insects, range from 40 to 75 g/100g and 10 to 30 g/100 g dry matter (DM), respectively (Verkerk et al., 2007; Raheem et al., 2019). Additionally, insects contain significant amounts of vitamins like pantothenic acid, riboflavin, biotin, and folic acid) and minerals, such as iron, copper, magnesium, phosphorus, selenium, manganese, and zinc (Eswaran et al., 2022; Mabelebele et al., 2022). However, increasing interest in entomophagy has raised concerns about the safety of insect-based products, mainly from a microbiological and chemical perspective (Imathiu, 2020). Insects can carry bacteria dangerous to humans and animals, thus potentially acting as vectors of foodborne pathogens. Therefore, when looking at the microbiology of edible insects, it is essential to consider that insects naturally carry microorganisms and may harbour a complex autochthonous microbial population (Marshall et al., 2016; Garofalo et al., 2019; Raheem et al., 2019).

Freezing is the most common method of killing insects, but it may not guarantee adequate microbiological quality (Grabowski and Klein, 2017). Therefore, it is necessary to heat treat or include another effective technological treatment in insect processing since removing their digestive tracts is impractical on a large scale for farmed edible insects. Moreover, the safety of processed insects varies depending on the species and the production process (Aguilar-Toalá et al., 2022; Pasini et al., 2022; Ververis et al., 2022; Gałęcki et al., 2023). From this point of view, the production process should include a heat treatment step that must be optimised with respect to the insect species and the intended use of the harvested insect biomass. For these reasons, investigating the microbiota of edible insects is essential to assessing and addressing potential health risks. Therefore, this study aims to study the effect of various killing methods and subsequent culinary treatments on the microbiological quality of edible insects. The procedures chosen for the experiment (boiling, freezing) were designed to simulate domestic and

industrial insect processing. For this, the holometabolous mealworm (*Tenebrio molitor*, Linnaeus, 1758) (TM) and the hemimetabolous Jamaican field cricket (*Gryllus assimilis*, Fabricius, 1775) (GA) were chosen as model organisms. Both species belong to the category of insects that are commercially farmed globally and are widely considered to be highly promising sources of food and feed (Dourado et al., 2020; Oliveira et al., 2024).

MATERIAL AND METHODS

Experimental insects

Insects were obtained from the Department of Zoology and Fisheries rearing facility, Czech University of Life Sciences Prague. Both species were maintained at $27 \pm 1^\circ\text{C}$ with a relative humidity of 40–50%, lightning regime 12:12 using the rack system. The house crickets were reared in plastic boxes (56 × 39 × 28 cm, IKEA, Prague, Czech Republic) equipped with egg trays (Schubert Partner, Prague, Czech Republic) and secured using a lid comprising approximately 80% of aluminium anti-insect mesh. Petri dishes containing feed (77.9% wheat, 17.6% soybean meal, 1.8% rapeseed oil, and 2.7% minerals and vitamin premix; particle size < 1 mm) were provided either in case of shortage or every 72 hours with unconsumed substrate removed if present. Petri dishes containing water gel (Oslavan, Náměšť nad Oslavou, Czech Republic) were also provided. The colonies were started with approximately 0.5 grams of 1-day-old pinheads that remained in the same container until harvest at 55 ± 2 days. The containers and Petri dishes were washed with hot water and detergent. The feed substrate was stored hermetically in a container at room temperature. The mealworm colonies were started with approximately 1000 three-week-old larvae, which were reared in the insect breeding trays (60 × 40 × 12.5 cm, Beekenkamp, Maasdijk, Netherlands) using a mixture of wheat bran and the substrate as mentioned above in 4:1 ratio. The water gel was provided directly to the substrate as a source of moisture on a daily basis. The mealworms were harvested when the first pupae occurred in the boxes. All trays were washed in hot tap water with detergent prior to the experiment. The wheat bran was also stored hermetically at room temperature. Before the harvest, the insects were starved for 24 hours.

Sample preparation

The insects were killed using two killing methods: freezing (-18°C for 24 hours) and blanching in hot water (100°C for 1 minute). Following this, the insects

underwent technological processing using the methods outlined in **Table 1**. Once cool, the samples were aseptically wrapped and refrigerated.

Table 1 Technological treatments and their conditions

microwave treatment	TM – 260 g for 10 min, GA 240 g for 10 min, 800 W
boiling	300 g for 30 min in 3 L boiling water
roasting	150 g, roasting without oil for 5 min
drying	60 g GA, 80 g TM, 15 hours, 80°C

Microbiological analysis

Sample preparation involved aseptically homogenised insect biomass by hand in a mixing bowl. One gram was then transferred to 9 ml of dilution medium (9 g of NaCl, 1 g of peptone (Oxoid, England), 1000 ml of distilled water) and serially diluted to a value of up to 10⁻⁹. All groups of microorganisms were identified and enumerated using classic culture methods in compliance with the following valid standards and procedures: for total aerobic spores, ISO 4833-1:2013 using Tryptone-Soy Agar (Oxoid, England); for *Bacillus cereus*, ISO 7932:2004 using *B. cereus* agar base (Himedia, USA) enriched with polymyxin B (2 vials/l, Himedia, USA) and yolk emulsion (50 ml/l, Himedia, USA); for *Escherichia coli*, ISO 16649-2:2001 with TBX medium (Oxoid, England); for *Enterobacteriaceae*, ISO 21528-2:2017 using Violet red bile glucose agar (Oxoid, England); for Coagulase-positive staphylococci and *Staphylococcus aureus*, ISO 6888-1:2021 using S Baird-Parker agar (Oxoid, England) with yolk emulsion and potassium tellurite; for *Salmonella* spp. ISO 6579-1:2017 using Rappaport-Vassiliadis (Oxoid, England) and SS agar (Oxoid, England); and for *Listeria monocytogenes* ISO 11290-1:2017 using Brilliance Listeria Agar (Oxoid, England). Coagulase activity was tested using the Staphylase test kit (Oxoid, England).

Statistical analysis

Statistical evaluation was performed using Statgraphics Centurion XV 15.2.05/2007 software (StatPoint Technologies, USA) employing one-way

analysis of variance (ANOVA) followed by Fisher's least significant difference as a post-hoc test, with a 95% confidence interval.

RESULTS

Results are presented in **Tables 2** and **3**. For both insect species, it is evident that the method of killing significantly affects the presence of the assessed groups of microorganisms. In GA, there is a significant decrease in aerobic sporulating bacteria from 5.78 ±0.03 log CFU/g in samples killed by freezing compared to 3.53 ±0.09 log CFU/g after blanching. A similar trend is observed for *B. cereus* and other sporulating bacteria. Values of aerobic sporulating bacteria after killing by blanching without other processing method were not significantly different to the values from all types of technological treatment. For TM, the most significant difference between killing methods was observed for *Enterobacteriaceae*, with 4.71 ±0.08 log CFU/g remaining after freezing compared to 1.48 ±0.09 log CFU/g after blanching. Moreover, a significant decrease in the total number of mesophilic aerobic bacteria when blanched was observed, while boiling and microwave treatment gave similar values to blanching (control samples).

For GA and TM, all types of treatment could eliminate *Enterobacteriaceae*, which could not be achieved by killing alone. Similar results apply to coagulase-positive staphylococci, while *L. monocytogenes*, *Salmonella* spp. and *E. coli* were not observed in the tested samples. The lowest counts for *B. cereus* in both TM and GA were observed in the boiled samples. However, exact log CFU values could not be determined in the samples due to recurrent overgrowth of the colonies over the whole surface of Petri dishes.

In TM, the lowest significant log CFU/g values for mesophilic aerobic bacteria were detected after roasting in insects killed by blanching. In GA, drying was the most effective method, regardless of the killing method. Roasted samples also produced the same values, but only for blanched insects, while in GA, the lowest significant values of sporulating aerobic bacteria were recorded after roasting frozen and blanched insects. Notably, all treatments significantly reduced the number of aerobic spore-formers in TM, although significantly higher values were recorded for boiled samples after freezing than other treatments.

Table 2 Results of microbiological indicators for *Gryllus assimilis* (log CFU/g). Values with different superscripts indicate significant difference (p < 0.05).

	control		drying		boiling		microwave treatment		roasting	
	freezing	blanching	freezing	blanching	freezing	blanching	freezing	blanching	freezing	blanching
<i>Bacillus cereus</i>	<4	<2	<2	<2	<1	<1	<2	<2	<2	<2
<i>Enterobacteriaceae</i>	5.85 ±0.09	4.75 ±0.10	<1	<1	<1	<1	<1	<1	<1	<1
<i>Escherichia coli</i>	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1
Coagulase-positive staphylococci	1.44 ±0.24	<1	<1	<1	<1	<1	<1	<1	<1	<1
<i>Staphylococcus aureus</i>	0	0	0	0	0	0	0	0	0	0
mesophilic aerobic bacteria	7.94 ±0.18 ^d	6.82 ±0.06 ^c	4.72 ±0.19 ^a	4.66 ±0.09 ^a	5.61 ±0.18 ^b	4.69 ±0.17 ^a	5.71 ±0.06 ^b	5.73 ±0.15 ^b	5.66 ±0.06 ^b	4.66 ±0.16 ^a
aerobic sporulating bacteria	5.78 ±0.03 ^d	3.53 ±0.09 ^{abc}	3.54 ±0.06 ^{abc}	3.58 ±0.11 ^{abc}	3.68 ±0.18 ^c	3.48 ±0.10 ^{ab}	3.62 ±0.07 ^{bc}	3.61 ±0.10 ^{bc}	3.44 ±0.05 ^a	3.46 ±0.09 ^{ab}
<i>Listeria monocytogenes</i>	0/25 g	0/25 g	0/25 g	0/25 g	0/25 g	0/25 g	0/25 g	0/25 g	0/25 g	0/25 g
<i>Salmonella</i>	0/25 g	0/25 g	0/25 g	0/25 g	0/25 g	0/25 g	0/25 g	0/25 g	0/25 g	0/25 g

Table 3 Results of microbiological indicators for *Tenebrio molitor* (log CFU/g). Values with different superscripts indicate significant difference (p < 0.05).

	control		drying		boiling		microwave treatment		roasting	
	freezing	blanching	freezing	blanching	freezing	blanching	freezing	blanching	freezing	blanching
<i>Bacillus cereus</i>	3.63 ±0.24	<2	<2	<2	<1	<1	<2	<2	<2	<2
<i>Enterobacteriaceae</i>	4.71 ±0.08	1.48 ±0.09	<1	<1	<1	<1	<1	<1	<1	<1
<i>Escherichia coli</i>	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1
Coagulase-positive staphylococci	1.33 ±0.15	1.35 ±0.24	<1	<1	<1	<1	<1	<1	<1	<1
<i>Staphylococcus aureus</i>	0	0	0	0	0	0	0	0	0	0
mesophilic aerobic bacteria	7.97 ±0.07 ^d	5.69 ±0.05 ^c	4.67 ±0.14 ^b	4.71 ±0.27 ^b	5.85 ±0.08 ^c	5.79 ±0.12 ^c	5.79 ±0.12 ^c	5.73 ±0.14 ^c	4.71 ±0.09 ^b	3.67 ±0.04 ^a
aerobic sporulating bacteria	5.76 ±0.03 ^c	4.68 ±0.13 ^b	3.59 ±0.15 ^a	3.62 ±0.06 ^a	4.51 ±0.06 ^b	3.63 ±0.14 ^a	3.56 ±0.15 ^a	3.62 ±0.11 ^a	3.62 ±0.21 ^a	3.54 ±0.04 ^a
<i>Listeria monocytogenes</i>	0/25 g	0/25 g	0/25 g	0/25 g	0/25 g	0/25 g	0/25 g	0/25 g	0/25 g	0/25 g
<i>Salmonella</i>	0/25 g	0/25 g	0/25 g	0/25 g	0/25 g	0/25 g	0/25 g	0/25 g	0/25 g	0/25 g

DISCUSSION

After comparing many studies, **Garafalo et al. (2019)** concluded that there were significant differences in the microbial load among different species of edible insects. Fresh edible insects generally contain a high microbial load, with mesophilic aerobes ranging from 3.6 to 9.4 log CFU/g, *Enterobacteriaceae* from 4.2–7.8 log CFU/g, and sporulating aerobic bacteria from 0.5–5.8 log CFU/g. In this study, the values for GA and TM also fall within these ranges. Moreover, the different microbial loads between TM and GA are evident, especially in the case of *Enterobacteriaceae*, where GA counts are two orders of magnitude higher than those of TM. TM is typically reared in a feed substrate of wheat bran, supplemented with fresh fruit and vegetables. In addition, they are in close contact with their frass. On the other hand, crickets are mainly fed a dry soy-based substrate and are not in close contact with their frass, which may explain the differences in the composition of the microorganisms. Notably, this study reveals a significant difference in the number of bacteria in insects killed by freezing and blanching. **Speck and Ray (1977)** stated that although some microorganisms may be killed during freezing, many survive in different states of viability. Increased resistance to freezing is usually provided by food components such as proteins, simple and complex carbohydrates, and triacylglycerols. Insects are rich in proteins (**Kouřimská et al., 2023**) and can serve as cryoprotectants, thereby protecting the microorganisms present, especially in the digestive tract, since insects are only starved before being killed. From a microbiological perspective, blanching appears more efficient, although, in the case of GA, there is a risk of insects escaping during handling.

After treatment, the lower microbial count is apparent compared to insects killed without any further treatment, which shows their effectiveness in reducing microbial contamination, agreeing with the findings of **Garafalo et al. (2019)**, **Aguilar-Toalá et al. (2022)**, **Pasini et al. (2022)**, **Ververis et al. (2022)**, and **Gałęcki et al. (2023)**. Furthermore, the present study found a significant decrease in microorganism counts after blanching. This finding agrees with similar results reported by **Vandeweyer et al. (2017)**, **Wynants et al. (2017)**, **Mancini et al. (2019)**, and **Cacchiarelli et al. (2022)**. **Nyangena et al. (2020)** reported roasting and drying in a hot air oven as the most effective methods for reducing the microbial load, boiling was also a suitable treatment. These findings confirm the results obtained for GA, which, regardless of the killing method, showed the lowest numbers of mesophilic bacteria after drying and TM, which had the lowest values after roasting. Similarly, boiling showed the greatest effect on reducing the amount of *Bacillus cereus*, i.e., by one order of magnitude compared to other types of treatment.

Although *L. monocytogenes*, *Salmonella* spp. and *E. coli* are important microbiological parameters in evaluating food quality, they were not detected in any of our samples, including untreated ones. Similar findings were reported by other studies, such as **Garafalo et al. (2019)**, **Kolakowski et al. (2021)**, and **Ververis et al. (2022)**.

The total number of aerobic bacteria represents one of the microbial guidelines used to assess the quality of fresh foods, with high total aerobic numbers being associated with rapid spoilage and potential health risks (**Ssepuuya et al., 2019**). Specifically, for the house cricket, data on the microbial quality of unprocessed samples showed a high load of total mesophilic bacteria ranging from 7.2 to 10.2 log CFU/g (**Ververis et al., 2022**). For GA, another cricket species, the results of this study produced similar values: 7.94 ± 0.18 log CFU/g. **Stoops et al. (2016)** found total aerobic numbers for unprocessed TM of 7.7–8.3 log CFU/g, agreeing with the present study's results (7.97 ± 0.07 log CFU/g). Similarly, **Klunder et al. (2012)** also reported total aerobic bacterial counts for fresh TM and house cricket in the 6.7–7.7 log CFU/g range. However, after technological modifications (boiling, roasting and frying), the total aerobic bacteria counts were <1.7–4.8 log CFU/g. In our case, the samples treated by roasting and drying did not exceed 4.8 log CFU/g, but for boiling and microwave radiation treatment, the total aerobic bacteria count was higher (TM: 5.73–5.85 and for GA: 4.69–5.73 log CFU/g). These findings indicate that the microwave treatment applied could not reduce total aerobic mesophilic bacteria after blanching.

Staphylococcus aureus is a common human pathogen present in edible insects. Contamination can be caused during handling or processing (**Garafalo et al., 2017**; **Milanović et al., 2018**; **Garafalo et al., 2019**). It is also known for producing toxins resistant to heat treatment, freezing, and drying (**Kooh et al., 2020**). However, *S. aureus* is sensitive to heat processing, and processed edible insects are favourable for the growth of this bacterium because of its ability to thrive without competition. Moreover, it is also resistant to low water activities (**Milanović et al., 2018**; **Walia et al., 2018**; **Garafalo et al., 2019**; **Kooh et al., 2019**). In our experiment, although the presence of *S. aureus* was not demonstrated, coagulase-positive staphylococci were detected. However, in TM and GA, they were destroyed by all treatment types, regardless of the killing method. However, in other studies, the presence of *S. aureus* was demonstrated in fresh and processed edible insects: yellow mealworm (**Garafalo et al., 2017**; **Wynants et al., 2017**; **Milanović et al., 2018**), lesser mealworm (**Wynants et al., 2018**), house cricket and desert locust (**Garafalo et al., 2017**; **Milanović et al., 2018**). Therefore, evaluating its presence in edible insects is necessary to ensure their safe use in the food industry.

With some exceptions, bacteria belonging to the family of *Enterobacteriaceae* are relatively sensitive to heat treatment (**Stoops et al., 2017**). For example, **Fröhling et al. (2020)** found that enterobacteria prevailed in untreated and treated house crickets and **Messina et al. (2019)** in house crickets and yellow mealworms. In the present study, untreated TM samples also had relatively high values of *Enterobacteriaceae* (4.71 ± 0.08 log CFU/g). Nonetheless, blanching alone resulted in a significant reduction of *Enterobacteriaceae* (1.48 ± 0.09 log CFU/g). For GA, the killing without further processing did not have such significant impact as for TM, resulting in values of 6.85 ± 0.09 to 4.75 ± 0.10 log CFU/g. **Osimani et al. (2017)** also found viable counts of *Enterobacteriaceae* (<1 log CFU/g) in dried house cricket and yellow mealworm, while **Klunder et al. (2012)** and **Stoops et al. (2017)** found that boiling for 10 minutes is sufficient to inactivate *Enterobacteriaceae*. Although the present study confirms these results, it also shows that all technological treatments can reduce *Enterobacteriaceae* to <1 log CFU/g in both species of edible insects and meet the specifications proposed by **EFSA (2021a, 2021b)**.

A potential risk associated with edible insects is the presence of spore-forming bacteria, which are likely introduced by contact with the soil, such as from the vegetables used as feed, which cannot be eliminated by boiling or drying (**Stoops et al., 2017**; **Walia et al., 2018**; **Garafalo et al., 2019**). **Fasolato et al. (2018)** and **Osimani and Aquilanti (2021)** also confirmed that drying and freeze-drying were ineffective against microbial spores. In our study, the most successful method of *B. cereus* reduction was processing by boiling. Also, blanching was a significantly more efficient killing method for reducing total aerobic spore count than freezing, but with further treatment (all tested treatments), a significant reduction in spore formers could be achieved. It has also been shown that heat treatment can reduce the number of bacterial spores and thus increase the possible shelf life, but at the expense of the nutritional and sensory quality of the final product (**Klunder et al., 2012**; **Grabowski and Klein, 2017**; **Vandeweyer et al., 2017**). **Osimani et al. (2017)** and **Fasolato et al. (2018)** pointed out that *B. cereus* was found across the species of edible insects. In our study, its presence was found in both two monitored species. *Bacillus cereus* is a foodborne pathogen, and a value of 5 log CFU/g was determined as microbial load that can cause gastrointestinal disease after ingestion (**Kramer and Gilbert, 1989**; **Garafalo et al., 2019**). **Osimani et al. (2017)** reported *B. cereus* counts above the critical level of 5 log CFU/g in processed insects, that are now commercially available in Europe, including house cricket meal samples. In the present study, untreated samples showed >5 log CFU/g in both species.

Based on the results of this study, we agree with the conclusions of **Ververis et al. (2022)**, who stated that technological processing is essential to ensure the microbiological safety of edible insects. Importantly, the quality of unprocessed insects needs to be monitored at least in terms of the following microbiological criteria: total numbers of mesophilic bacteria, spore-forming bacteria, *B. cereus*, *S. aureus*, *Enterobacteriaceae* and coagulase-positive staphylococci. These microorganisms were the most prevalent in untreated samples of TM and GA in our study.

CONCLUSION

Based on the analyses conducted in this study, the presence of *B. cereus*, mesophilic aerobic bacteria, aerobic sporulating bacteria and bacteria of the genus *Enterobacteriaceae* were detected in the samples GA and TM. Pathogenic bacteria such as *S. aureus*, *Salmonella* spp., and *L. monocytogenes* remained undetected in any of samples. After the killing of the edible insects, the microbial load remained relatively high. Killing by blanching in hot water helps to reduce the microbial load more than by freezing, but even so, additional culinary treatment did show an effect and is recommended for consumer safety to prevent potential exposure to pathogenic bacteria. In summary, insects killed by blanching or freezing without further technological processing should not be considered microbiologically safe for consumption.

Acknowledgements: The authors thank Dora Petříčková, Kristýna Horváthová, Marek Hauk, Petra Belfínová and Alice Andrlová for their help with sample preparation and laboratory analysis, and Michal Kurečka for maintaining the insect colonies in the insectarium. This project was supported by the Grant Agency of the Czech Republic GAČR project No. 21- 47159L and Slovenian Research Agency (ARIS, program group P1-0143 and project number J7-3155), bilateral project: Quality, safety and authenticity of insect protein-based food and feed products (INPROFF) and by the METROFOOD-CZ research infrastructure project (MEYS Grant No: LM2023064) including access to its facilities.

REFERENCES

- Aguilar-Toalá, J. E., Cruz-Monterrosa, R. G., & Liceaga, A. M. (2022). Beyond Human Nutrition of Edible Insects: Health Benefits and Safety Aspects. *Insects*, 13(11). <https://doi.org/10.3390/insects13111007>
- Cacchiarelli, C., Fratini, F., Puccini, M., Vitolo, S., Paci, G., & Mancini, S. (2022). Effects of different blanching treatments on colour and microbiological profile of

- Tenebrio molitor* and *Zophobas morio* larvae. *LWT*, 157. <https://doi.org/10.1016/j.lwt.2022.113112>
- Dourado, L. R. B., Lopes, P. M., Silva, V. K., Carvalho, F. L. A., Moura, F. A. S., Silva, L. B., Giannecchini, L. G., Pinheiro, S. R. F., Biagiotti, D., & Kimpara, J. M. (2020). Chemical composition and nutrient digestibility of insect meal for broiler. *Anais da Academia Brasileira de Ciências*, 92(3). <https://doi.org/10.1590/0001-37652020200764>
- EFSAA. (2021). Safety of frozen and dried formulations from whole yellow mealworm (*Tenebrio molitor* larva) as a novel food pursuant to Regulation (EU) 2015/2283. *EFSA Journal*, 19(8). <https://doi.org/10.2903/j.efsa.2021.6778>
- EFSAB. (2021). Safety of frozen and dried formulations from whole house crickets (*Acheta domestica*) as a Novel food pursuant to Regulation (EU) 2015/2283. *EFSA Journal*, 19(8). <https://doi.org/10.2903/j.efsa.2021.6779>
- Eswaran, U. G. M., Karunanithi, S., Gupta, R. K., Rout, S., & Srivastav, P. P. (2022). Edible insects as emerging food products—processing and product development perspective. *Journal of Food Science and Technology*. <https://doi.org/10.1007/s13197-022-05489-y>
- Fasolato, L., Cardazzo, B., Carraro, L., Fontana, F., Novelli, E., & Balzan, S. (2018). Edible processed insects from e-commerce: Food safety with a focus on the *Bacillus cereus* group. *Food Microbiology*, 76, 296-303. <https://doi.org/10.1016/j.fm.2018.06.008>
- Fröhling, A., Büßler, S., Durek, J., & Schlüter, O. K. (2020). Thermal Impact on the Culturable Microbial Diversity Along the Processing Chain of Flour from Crickets (*Acheta domestica*). *Frontiers in Microbiology*, 11. <https://doi.org/10.3389/fmicb.2020.00884>
- Gałęcki, R., Bakula, T., & Gołaszewski, J. (2023). Foodborne Diseases in the Edible Insect Industry in Europe—New Challenges and Old Problems. *Foods*, 12(4). <https://doi.org/10.3390/foods12040770>
- Garofalo, C., Osimani, A., Milanović, V., Taccari, M., Cardinali, F., Aquilanti, L., Riolo, P., Ruschioni, S., Isidoro, N., & Clementi, F. (2017). The microbiota of marketed processed edible insects as revealed by high-throughput sequencing. *Food Microbiology*, 62, 15-22. <https://doi.org/10.1016/j.fm.2016.09.012>
- Garofalo, C., Milanović, V., Cardinali, F., Aquilanti, L., Clementi, F., & Osimani, A. (2019). Current knowledge on the microbiota of edible insects intended for human consumption: A state-of-the-art review. *Food Research International*, 125. <https://doi.org/10.1016/j.foodres.2019.108527>
- Grabowski, N. T., & Klein, G. (2017). Microbiology of processed edible insect products – Results of a preliminary survey. *International Journal of Food Microbiology*, 243, 103-107. <https://doi.org/10.1016/j.ijfoodmicro.2016.11.005>
- Imathiu, S. (2020). Benefits and food safety concerns associated with consumption of edible insects. *NFS Journal*, 18, 1-11. <https://doi.org/10.1016/j.nfs.2019.11.002>
- Klunder, H. C., Wolkers-Rooijackers, J., Korpela, J. M., & Nout, M. J. R. (2012). Microbiological aspects of processing and storage of edible insects. *Food Control*, 26(2), 628-631. <https://doi.org/10.1016/j.foodcont.2012.02.013>
- Kolakowski, B. M., Johaniuk, K., Zhang, H., & Yamamoto, E. (2021). Analysis of Microbiological and Chemical Hazards in Edible Insects Available to Canadian Consumers. *Journal of Food Protection*, 84(9), 1575-1581. <https://doi.org/10.4315/JFP-21-099>
- Kooh, P., Ververis, E., Tesson, V., Boué, G., & Federighi, M. (2019). Entomophagy and Public Health: A Review of Microbiological Hazards. *Health*, 11(10), 1272-1290. <https://doi.org/10.4236/health.2019.1110098>
- Kooh, P., Jury, V., Laurent, S., Audiat-Perrin, F., Sanaa, M., Tesson, V., Federighi, M., & Boué, G. (2020). Control of Biological Hazards in Insect Processing: Application of HACCP Method for Yellow Mealworm (*Tenebrio molitor*) Powders. *Foods*, 9(11). <https://doi.org/10.3390/foods9111528>
- Kouřimská, L., Kvasnička, F., Kurečka, M., Rajchl, A., Škvorová, P., & Kulma, M. (2023). Taurine content of insects used as feed. *Journal of Insects as Food and Feed*, 9(9), 1151 - 1157. <https://doi.org/10.3920/JIFF2022.0140>
- Kramer, J. M., & Gilbert, R. J. (1989). *Bacillus cereus* and other *Bacillus* species. *Foodborne bacterial pathogens*, 19: 21-70.
- Mabelebele, M., Kolobe, S. D., Malematja, E., Sebola, N. A., & Manyelo, T. G. (2022). A Comprehensive Review of the Importance of Selected Trace Elements Present in Edible Insects. *Biological Trace Element Research*. <https://doi.org/10.1007/s12011-022-03423-z>
- Mancini, S., Fratini, F., Tuccinardi, T., Turchi, B., Nuvoloni, R., & Paci, G. (2019). Effects of different blanching treatments on microbiological profile and quality of the mealworm (*Tenebrio molitor*). *Journal of Insects as Food and Feed*, 5(3), 225-234. <https://doi.org/10.3920/JIFF2018.0034>
- Marshall, D. L., Dickson, J. S., & Nguyen, N. H. (2016). Ensuring Food Safety in Insect Based Foods: Mitigating Microbiological and Other Foodborne Hazards. *Insects as Sustainable Food Ingredients* (223-253). Elsevier. <https://doi.org/10.1016/B978-0-12-802856-8.00008-9>
- Messina, C. M., Gaglio, R., Morghese, M., Tolone, M., Arena, R., Moschetti, G., Santulli, A., Francesca, N., & Settanni, L. (2019). Microbiological Profile and Bioactive Properties of Insect Powders Used in Food and Feed Formulations. *Foods*, 8(9). <https://doi.org/10.3390/foods8090400>
- Milanović, V., Osimani, A., Roncolini, A., Garofalo, C., Aquilanti, L., Pasquini, M., Tavoletti, S., Vignaroli, C., Canonico, L., Ciani, M., & Clementi, F. (2018). Investigation of the Dominant Microbiota in Ready-to-Eat Grasshoppers and Mealworms and Quantification of Carbenem Resistance Genes by qPCR. *Frontiers in Microbiology*, 9. <https://doi.org/10.3389/fmicb.2018.03036>
- Nyangena, D. N., Mutungi, C., Imathiu, S., Kinyuru, J., Affognon, H., Ekisi, S., Nakimbugwe, D., & Fiaboe, K. K. M. (2020). Effects of Traditional Processing Techniques on the Nutritional and Microbiological Quality of Four Edible Insect Species Used for Food and Feed in East Africa. *Foods*, 9(5). <https://doi.org/10.3390/foods9050574>
- Oliveira, L. A., Pereira, S. M. S., Dias, K. A., Paes, S. da S., Grancieri, M., Jimenez, L. G. S., de Carvalho, C. W. P., de Oliveira, E. E., Martino, H. S. D., & Della Lucia, C. M. (2024). Nutritional content, amino acid profile, and protein properties of edible insects (*Tenebrio molitor* and *Gryllus assimilis*) powders at different stages of development. *Journal of Food Composition and Analysis*, 125. <https://doi.org/10.1016/j.jfca.2023.105804>
- Oonincx, D. G. A. B., & Finke, M. D. (2021). Nutritional value of insects and ways to manipulate their composition. *Journal of Insects as Food and Feed*, 7(5), 639-659. <https://doi.org/10.3920/JIFF2020.0050>
- Osimani, A., Garofalo, C., Milanović, V., Taccari, M., Cardinali, F., Aquilanti, L., Pasquini, M., Mozzon, M., Raffaelli, N., Ruschioni, S., Riolo, P., Isidoro, N., & Clementi, F. (2017). Insight into the proximate composition and microbial diversity of edible insects marketed in the European Union. *European Food Research and Technology*, 243(7), 1157-1171. <https://doi.org/10.1007/s00217-016-2828-4>
- Osimani, A., & Aquilanti, L. (2021). Spore-forming bacteria in insect-based foods. *Current Opinion in Food Science*, 37, 112-117. <https://doi.org/10.1016/j.cofs.2020.10.011>
- Pasini, G., Cullere, M., Vegro, M., Simonato, B., & Dalle Zotte, A. (2022). Potentiality of protein fractions from the house cricket (*Acheta domestica*) and yellow mealworm (*Tenebrio molitor*) for pasta formulation. *LWT*, 164. <https://doi.org/10.1016/j.lwt.2022.113638>
- Raheem, D., Raposo, A., Oluwole, O. B., Nieuwland, M., Saraiva, A., & Carrascosa, C. (2019). Entomophagy: Nutritional, ecological, safety and legislation aspects. *Food Research International*, 126. <https://doi.org/10.1016/j.foodres.2019.108672>
- Shah, A. A., Totakul, P., Matra, M., Cherdthong, A., Harnboonsong, Y., & Wanapat, M. (2022). Nutritional composition of various insects and potential uses as alternative protein sources in animal diets. *Animal Bioscience*, 35(2), 317-331. <https://doi.org/10.5713/ab.21.0447>
- Speck, M. L., & Ray, B. (1977). Effects of Freezing and Storage on Microorganisms in Frozen Foods: A Review. *Journal of Food Protection*, 40(5), 333-336. <https://doi.org/10.4315/0362-028X-40.5.333>
- Ssepuya, G., Wynants, E., Verreth, C., Crauwels, S., Lievens, B., Claes, J., Nakimbugwe, D., & Van Campenhout, L. (2019). Microbial characterisation of the edible grasshopper *Ruspolia differens* in raw condition after wild-harvesting in Uganda. *Food Microbiology*, 77, 106-117. <https://doi.org/10.1016/j.fm.2018.09.005>
- Stoops, J., Vandeweyer, D., Crauwels, S., Verreth, C., Boeckx, H., Van Der Borcht, M., Claes, J., Lievens, B., & Van Campenhout, L. (2017). Minced meat-like products from mealworm larvae (*Tenebrio molitor* and *Alphitobius diaperinus*): microbial dynamics during production and storage. *Innovative Food Science & Emerging Technologies*, 41, 1-9. <https://doi.org/10.1016/j.ifset.2017.02.001>
- Vandeweyer, D., Lenaerts, S., Callens, A., & Van Campenhout, L. (2017). Effect of blanching followed by refrigerated storage or industrial microwave drying on the microbial load of yellow mealworm larvae (*Tenebrio molitor*). *Food Control*, 71, 311-314. <https://doi.org/10.1016/j.foodcont.2016.07.011>
- van Huis, A., & Oonincx, D. G. A. B. (2017). The environmental sustainability of insects as food and feed. A review. *Agronomy for Sustainable Development*, 37(5). <https://doi.org/10.1007/s13593-017-0452-8>
- van Huis, A. (2021). Prospects of insects as food and feed. *Organic Agriculture*, 11(2), 301-308. <https://doi.org/10.1007/s13165-020-00290-7>
- Verkerk, M. C., Tramper, J., van Trijp, J. C. M., & Martens, D. E. (2007). Insect cells for human food. *Biotechnology Advances*, 25(2), 198-202. <https://doi.org/10.1016/j.biotechadv.2006.11.004>
- Ververis, E., Boué, G., Poulsen, M., Pires, S. M., Niforou, A., Thomsen, S. T., Tesson, V., Federighi, M., & Naska, A. (2022). A systematic review of the nutrient composition, microbiological and toxicological profile of *Acheta domestica* (house cricket). *Journal of Food Composition and Analysis*, 114. <https://doi.org/10.1016/j.jfca.2022.104859>
- Walia, K., Kapoor, A., & Farber, J. M. (2018). Qualitative risk assessment of cricket powder to be used to treat undernutrition in infants and children in Cambodia. *Food Control*, 92, 169-182. <https://doi.org/10.1016/j.foodcont.2018.04.047>
- Wynants, E., Crauwels, S., Lievens, B., Luca, S., Claes, J., Borremans, A., Bruyninckx, L., & Van Campenhout, L. (2017). Effect of post-harvest starvation and rinsing on the microbial numbers and the bacterial community composition of mealworm larvae (*Tenebrio molitor*). *Innovative Food Science & Emerging Technologies*, 42, 8-15. <https://doi.org/10.1016/j.ifset.2017.06.004>
- Wynants, E., Crauwels, S., Verreth, C., Gianotten, N., Lievens, B., Claes, J., & Van Campenhout, L. (2018). Microbial dynamics during production of lesser mealworms (*Alphitobius diaperinus*) for human consumption at industrial scale. *Food Microbiology*, 70, 181-191. <https://doi.org/10.1016/j.fm.2017.09.012>